

2016-17 School Year New York State Immunization Requirements for School Entrance/Attendance¹

NOTES:

Children in a prekindergarten setting should be age-appropriately immunized. The number of doses depends on the schedule recommended by the Advisory Committee on Immunization Practices (ACIP). For grades Pre-k through 8, intervals between doses of vaccine should be in accordance with the ACIP-recommended immunization schedule for persons 0 through 18 years of age. (Exception: intervals between doses of polio vaccine need to be reviewed only for grades prekindergarten, kindergarten, 1, 2, 6, 7 and 8.) Doses received before the minimum age or intervals are not valid and do not count toward the number of doses listed below. Intervals between doses of vaccine DO NOT need to be reviewed for grades 9 through 12. See footnotes for specific information for each vaccine. Children who are enrolling in grade-less classes should meet the immunization requirements of the grades for which they are age equivalent.

Dose requirements MUST be read with the footnotes of this schedule.

Vaccines	Prekindergarten (Day Care, Head Start, Nursery or Pre-k)	Kindergarten and Grades 1 and 2	Grades 3, 4 and 5	Grades 6, 7 and 8	Grades 9, 10, 11 and 12
Diphtheria and Tetanus toxoid-containing vaccine and Pertussis vaccine (DTaP/DTP/Tdap) ²	4 doses	5 doses or 4 doses if the 4th dose was received at 4 years of age or older or 3 doses if aged 7 years or older and the series was started at 1 year of age or older		3 doses	
Tetanus and Diphtheria toxoid-containing vaccine and Pertussis vaccine booster (Tdap) ³		Not applicable		1 dose	
Polio vaccine (IPV/OPV) ⁴	3 doses	4 doses or 3 doses if the 3rd dose was received at 4 years of age or older	3 doses	4 doses or 3 doses if the 3rd dose was received at 4 years of age or older	3 doses
Measles, Mumps and Rubella vaccine (MMR) ⁵	1 dose		2 doses		
Hepatitis B vaccine ⁶	3 doses		3 doses or 2 doses of adult hepatitis B vaccine (Recombivax) for children who received the doses at least 4 months apart between the ages of 11 through 15 years of age		
Varicella (Chickenpox) vaccine ⁷	1 dose	2 doses	1 dose	2 doses	1 dose
Meningococcal conjugate vaccine (MenACWY) ⁸		Not applicable		By Grade 7: 1 dose	Grade 12: 2 doses or 1 dose if the dose was received at 16 years of age or older
Haemophilus influenzae type b conjugate vaccine (Hib) ⁹	1 to 4 doses		Not applicable		
Pneumococcal Conjugate vaccine (PCV) ¹⁰	1 to 4 doses		Not applicable		

1. Demonstrated serologic evidence of measles, mumps, rubella, hepatitis B, varicella or polio (for all three serotypes) antibodies is acceptable proof of immunity to these diseases. Diagnosis by a physician, physician assistant or nurse practitioner that a child has had varicella disease is acceptable proof of immunity to varicella.
2. Diphtheria and tetanus toxoids and acellular pertussis (DTaP) vaccine. (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive a 5-dose series of DTaP vaccine at ages 2 months, 4 months, 6 months and at 15 through 18 months and at 4 years of age or older. The fourth dose may be received as early as age 12 months, provided at least 6 months have elapsed since the third dose. However, the fourth dose of DTaP need not be repeated if it was administered at least 4 months after the third dose of DTaP. The final dose in the series must be received on or after the fourth birthday.
 - b. If the fourth dose of DTaP was administered at age 4 years or older, the fifth (booster) dose of DTaP vaccine is not required.
 - c. For children born before 1/1/2005, only immunity to diphtheria is required and doses of DT and Td can meet this requirement.
 - d. Children ages 7 through 10 years who are not fully immunized with the childhood DTaP vaccine series should receive Tdap vaccine as the first dose in the catch-up series; if additional doses are needed, use Td vaccine. A Tdap vaccine (or incorrectly administered DTaP vaccine) received at 7 years or age or older will meet the 6th grade Tdap requirement.
 - e. For children 7 years of age or older who received the first dose on or after their first birthday, the immunization requirement is 3 doses. If the first dose was received before their first birthday, then 4 doses are required.
3. Tetanus and diphtheria toxoids and acellular pertussis (Tdap) vaccine. (Minimum age: 7 years)
 - a. Students 11 years of age or older entering grades 6 through 12 are required to have one dose of Tdap. A dose received at 7 years of age or older will meet this requirement.
 - b. Students who are 10 years old in grade 6 and who have not yet received a Tdap vaccine are in compliance until they turn 11 years of age.
4. Poliovirus vaccine (IPV/OPV). (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive a series of IPV at ages 2 months, 4 months and at 6 through 18 months, and 4 years of age or older. The final dose in the series must be received on or after the fourth birthday and at least 6 months after the previous dose.
 - b. For students who received their fourth dose before age 4 and prior to August 7, 2010, 4 doses separated by at least 4 weeks is sufficient.
 - c. If the third dose of polio vaccine was received at age 4 years or older and at least 6 months after the previous dose, the fourth dose of polio vaccine is not required.
5. Measles, mumps, and rubella (MMR) vaccine. (Minimum age: 12 months)
 - a. The first dose of MMR vaccine must have been received on or after the first birthday. The second dose must have been received at least 28 days (4 weeks) after the first dose to be considered valid.
 - b. Students in grades kindergarten through 12 must have received 2 doses of measles-containing vaccine, 2 doses of mumps-containing vaccine and at least 1 dose of rubella-containing vaccine.
 - c. One dose of MMR is required for prekindergarten.
6. Hepatitis B vaccine
 - a. Dose 1 may be given at birth or anytime thereafter. Dose 2 must be given at least 4 weeks (28 days) after dose 1. Dose 3 must be at least 8 weeks after dose 2 AND at least 16 weeks after dose 1 AND no earlier than 24 weeks of age.
 - b. Two doses of adult hepatitis B vaccine (Recombivax) received at least 4 months apart at age 11 through 15 years will meet the requirement.
7. Varicella (chickenpox) vaccine. (Minimum age: 12 months)
 - a. The first dose of varicella vaccine must have been received on or after the first birthday. The second dose must have been received at least 28 days (4 weeks) after the first dose to be considered valid.
 - b. For children aged less than 13 years, the recommended minimum interval between doses is 3 months (if the second dose was administered at least 4 weeks after the first dose, it can be accepted as valid); for persons aged 13 years and older, the minimum interval between doses is 4 weeks.
8. Meningococcal conjugate vaccine (MenACWY). (Minimum age: 6 weeks)
 - a. One dose of meningococcal conjugate vaccine (Menactra or Menevo) is required for students entering grade 7.
 - b. For students in grade 12, if the first dose of meningococcal conjugate vaccine was received at age 16 years or older, the second (booster) dose is not required.
9. Haemophilus influenzae type b (Hib) conjugate vaccine. (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive Hib vaccine at 2 months, 4 months, 6 months and at 12 through 15 months. Children older than 15 months must get caught up according to the ACIP catch-up schedule. The final dose must be received on or after 12 months of age.
 - b. If 2 doses of vaccine were received before 12 months of age, only 3 doses are required with dose 3 at 12 through 15 months of age and at least 8 weeks after dose 2.
 - c. If dose 1 was received at ages 12 through 14 months of age, only 2 doses are required with dose 2 at least 8 weeks after dose 1.
 - d. If dose 1 was received at 15 months of age or older, only 1 dose is required.
 - e. Hib vaccine is not required for children 5 years of age or older.
10. Pneumococcal conjugate vaccine (PCV). (Minimum age: 6 weeks)
 - a. Children starting the series on time should receive PCV vaccine at 2 months, 4 months, 6 months and at 12 through 15 months. Children older than 15 months must get caught up according to the ACIP catch-up schedule. The final dose must be received on or after 12 months of age.
 - b. Unvaccinated children 7 through 11 months of age are required to receive 2 doses, at least 4 weeks apart, followed by a third dose at age 12 through 15 months.
 - c. Unvaccinated children 12 through 23 months of age are required to receive 2 doses of vaccine at least 8 weeks apart.
 - d. If one dose of vaccine was received at 24 months of age or older, no further doses are required.
 - e. For further information, refer to the PCV chart available in the School Survey Instruction Booklet at: www.health.ny.gov/prevention/immunization/schools

For further information contact:

New York State Department of Health
 Bureau of Immunization
 Room 649, Corning Tower ESP
 Albany, NY 12237
 (518) 473-4437

New York City Department of Health and Mental Hygiene
 Program Support Unit, Bureau of Immunization,
 42-09 28th Street, 5th floor
 Long Island City, NY 11101
 (347) 396-2433



Dear Parents:

All students entering **7th** and **12th** grade must have the **meningococcal vaccine**.

Without it, they cannot start school!

About the Vaccine:

- It's not a new vaccine. It's been recommended for a decade.
- Most parents already choose to vaccinate their children.

About Meningococcal Disease:

- It causes **bacterial meningitis** and other serious diseases.
- Teens and young adults are at a greater risk.
- It comes on quickly and without warning.
- Its symptoms are similar to the flu.
- Every case of this disease can result in death or long term disability.

Check with your doctor. Even kids who have had a shot before may need a booster to start school. Immunization is Protection

To learn more, visit: www.health.ny.gov/immunize.

Before school this September:

- **One dose of meningococcal vaccine is required before 7th grade.** If your child had the first dose as a sixth grader, then another dose is not required until 12th grade.
- **A total of 2 doses will be required before 12th grade.** Most students entering 12th grade got their first dose when they were younger and are now due for their second dose, or booster. This booster is needed because protection from the vaccine decreases over time.
- The **only** teens who will not need a 2nd dose before 12th grade are those who got their first dose on or after their 16th birthday.

It's best to check with your doctor to see whether or not your child needs the vaccine. Students who are not up-to-date will not be allowed to attend school until they are vaccinated.

Please call Nurse Bentley with any questions at 283-2500, ext. 235.